Birds of the North and North-West of Australia.

From Notes and Skins made by the late Capt. T. H. Bowyer-Bower.

By Gregory M. Mathews, F.R.S.E., F.L.S., M.B.O.U., etc. No. 12

156. CACOMANTIS PYRROPHANUS DUMETORUM. Western Square-tailed Cuckoo.

Cuculus dumetorum (Gould), Proc. Zool. Soc. (Lond.), 1845, p. 19. Port Essington, Northern Territory,

No. 308-9. Length, 8.5 inches (12/9/86).

Irides, brown round the pupil, and gradually shading into olive grey on its outer circle; upper mandible, black, lower, olive-brown; front of tarsi and feet brownish ochre; back of tarsi and soles of feet, dirty yellow; inside of bill and mouth, bright salmon colour, deepest on the latter.

No. 348— 3. Length, 9.0 inches (19/9/86).

.No. 445— 5. Length, 8.9 inches (26/10/86).

No. 527—3. Length, 9.0 inches (8/11/86).

156. Owenavis osculans rogersi. Western Black-eared Cuckoo.

Owenavis osculans rogersi (Mathews), Austral. Av. Rec., Irides, brown, bill and feet, black.

Vol. I., p. 13, 1912. Parry's Creek, North-west Australia.

No. 218—9. Length, 7.85 inches (5/7/86). No. 358—9. Length, 8.1 inches (2/8/86).

No. 359—3. Length, 7.8 inches (2/8/86).

No. 411—9. Length, 8.1 inches (18/10/86).

These birds were observed in a thick tea-tree scrub.

157. NEOCHALCITES BASALIS WYNDHAMI. Western Narrowbilled Bronze Cuckoo.

Chrysococcyx basalis wyndhami (Mathews), Austral. Av. Rec., Vol. I., p. 14, 1912. Point Torment, North-west Australia.

No. 118—Length, 6.8 inches (12/6/86).

Irides, brownish grey; bill, black, bluish at the base of the lower mandible; feet mealy black.

160. Polophilus phasianinus melanurus. North-western

Centropus melanurus (Gould), Birds, Austral., Vol. IV., text te pl. 92, 1847. Derby, North-west Australia.

No. 202— 9. Length, 2 feet, 3 inches (28/6/86).

No. $214 - 9 \cdot (1/7/86)$.

Irides, light greenish brown; space around the eye, bluish lead colour; bill, fleshy white; legs and feet, lead colour.

No. 220— & Length, 25.7 (26/8/86).

165. Hylochelidon nigricans neglecta. Western Tree Martin.

. Petrochelidon nigricans neglecta (Mathews), Nov. Zool., Vol. XVIII., p. 301, 1912. Fitzroy River, North-west Australia.

No. 297— & . Length, 4.8 inches.

No. 298.—Length, 4.7 inches (21/7/86).

Bill and legs, black. Very common, occasionally rests on dead trees in considerable numbers, chiefly in the early morning.

166. MICROECA FASCINANS SUBPALLIDA. Western Pale Flycatcher.

Microeca fascinans subpallida (Mathews), Nov. Zool., Vol. XVIII., p. 302, 1912. Napier Broome Bay, North-west Australia.

No. 54—9. Length, 5.2 inches (27/5/86).

Irides, dark brown; bill, brown; legs, black.

No. 83—9. Length, 5.1 inches (2/6/86).

No. 84— δ . Length, 5.1 inches (2/6/86).

Irides, dark brown; bill, black, whitish at the base of the lower mandible; feet black.

170. MELANODRYAS CUCULLATA PICATA. Pied Robin.

Melanodryas picata (Gould), Handb. Birds Austral., Vol. I., p. 285, 1865. North-west Coast, Australia.

No. 487— &. Length, 6.5 inches.

No. 492—9. Length, 5.7 (2/11/86).

Irides, dark brown; bill, legs, and feet, black.

No. 499—6. Length, 6.4 inches (3/11/06).

No. 502—3. Length, 6.4 inches (3/11/06).

Irides, brown; bill, legs and feet, black.

No. 518-9. Length, 6.2 inches (5/11/86).

171. SMICRORNIS BREVIROSTRIS ROGERSI. North-western Tree

Smicrornis brevirostris rogersi (Mathews), Nov. Zool., Vol. XVIII., p. 307, 1912. Napier Broome Bay, North-west Australia.

No. 191— 3. Length, 3.2 (27/6/06).

No. 193— \circ . Length, 3.2 27/6/06).

No. 209—9. Length, 3.5 (29/6/06). No. 291—3. Length, 3.4 (19/7/86).

No. 352—Length, 3.2 (2/8/86).

No. 353—Length, 3.4 (2/8/86).

No. 368—Length, 3.7 (26/9/86).

172. Gerygone olivacea rogersi. Grey Fly-eater.

Gerygone albigularis rogersi (Mathews), Nov. Zool., Vol.

XVIII., p. 23, 1911. Derby, North-west Australia. No. 95—3. Length 4 inches (5/6/86).

Bill, brown, fleshy white at the base of the lower mandible; legs, bluish-grey.

No. 113— &. Length, 4.1 inches (8/6/86).

Irides, light red; bill, dark brown; feet, dark grey.

No. 115-9. Length, 4 inches (8/6/86).

Bill, dark brown, lighter at the base of the lower mandible.

No. 275— δ . Length, 4 inches (5/9/86).

176. Poecilodryas superciliosa belcheri. Western Buffsided Robin.

Pachycephala superciliosa belcheri (Mathews), Austral., Av. Rec. Vol. I., p. 40, 1912. Napier Broome Bay, North-west Australia.

No. 221—3. Length, 6.1 inches (27/8/86).

No. 267— δ . Length, 6.1 inches (3/9/86).

No. 305— 3. (11/9/06).

No. 330—Length, 6.9 (16/9/86).

No. 332—Length 7 inches (16/9/86).

Irides, brown; feet and legs, brown.

No. 337 - 9. Length, 6.4 inches 17/9/86).

Irides, greyish brown; bill, black; feet and legs, brown.

No. 465—imm. Length, 4.7 inches (28/10/86).

Irides, pale slaty brown; the pupil with a bluish and dull tint; eyelid, brown; somewhat lighter than the feathers of the head; bill, brown; the cutting edges and gape yellowish white; legs and feet light brown, with a purplish tint on the tarsus; claws, white.

No. 467— & . Length, 6.8 (28/10/86).

No. 468— 9. Length, 6.5 (28/10/86).

Irides, brown; bill, feet, and legs, black.

The song of this bird is most beautiful and varied. In itself also it is a lovely bird when alive. The skins give no idea of its real beauty, when seen sitting under the shade of a tree giving forth its beautiful song, with the dark green as a background. It is very robin-like in its movements, and when singing frequently jerks up its head after the manner of that bird. It moves along the branches in a very taking manner, and when singing lets its wings fall below the tail; the colours show out very brightly.

The two adults and young were obtained in a very shady spot. On the male being wounded and falling to the ground, the female went down to him, and both were secured, a pair not having been seen together before. The young one could fly, and also had to be shot.

178. Раснусернава PECTORALIS MELANURA, Black-tailed Thickhead.

Pachycephala melanura (Gould), Proc. Zool. Soc. (Lond.), 1842, p. 134, 1843. Derby, North-west Australia.

No. 311— &. Length, 6.6 inches (25/7/86).

180. Lewinornis rufiventris colletti. Western Rufousbreasted Thickhead.

Pachycephala rufiventris colletti (Mathews), Austral. Av. .Rec., Vol. I., p. 41. Parry's Creek, North-west Australia.

No. 188— & Length, 6.5 inches (26/6/86).

Irides, reddish brown; bill and feet, black.

No. 189-9. Length, 6.2 (26/6/86).

Bill brown, lower mandible lighter at the base. The above are a pair shot together.

No. 197— & Length, 6.7 (27/6/86).

Irides, reddish brown; bill, legs, and feet, black.

No. 275—9. Length, 6.8 inches (17/7/86). No. 354—3. Length, 6.7 inches (2/8/86).

No. 476— δ . Length, 6.7 inches (29/10/86)

The males of this subspecies are very shy, and not easily obtained, the females and young birds are easily shot. note is full and sweet.