

*Bird Notes.*

By H. Simpson, Edward's Creek, S.A.

Edward's Creek is 624 miles north of Adelaide, and 64 miles south of Oodnadatta, by rail. The following species have been observed in the district:—Diamond Dove (*Geopelia cuneata*);

*Bird Notes.*

Crested Pigeon (*Ocyphaps lophotes*); Southern Stone Curlew (*Burhinus magnirostris*); Bustard (*Eupodotis australis*); Grey Teal (*Querquedula gibberifrons*); Wedge-tailed Eagle (*Uroaetus audax*); Black-shouldered Kite (*Elanus axillaris*); Brown Hawk (*Falco berigora*); Nankeen Kestrel (*Falco cenchroides*); White Cockatoo (*Kakatoe galerita*) (?); Galah (*K. roseicapilla*); Cockatoo Parrot (*Leptolophus hollandicus*); Port Lincoln Parrot (*Barnardius zonarius*); Mulga Parrot (*Psephotus varius*); Budgerygah (*Melopsittacus undulatus*); Owlet Nightjar (*Aegotheles cristata*); Bee-eater (*Merops ornatus*); Welcome Swallow (*Hirundo neoxena*); Tree-Martin (*Hylochelidon nigricans*); Fairy Martin (*H. ariel*); Magpie-Lark (*Grallina cyanoleuca*); Crested Bellbird (*Oreoica gutturalis*); Black-faced Cuckoo-Shrike (*Coracina novae-hollandiae*); Brown Songlark (*Cinclorhamphus cruralis*); Blue and White Wren (*Malurus cyanotus*); White-plumed Honeyeater (*Meliphaga penicillata*); Yellow-throated Miner (*Myzantha flavigula*); Spiny-cheeked Honeyeater (*Acanthagenys rufogularis*); Chestnut-eared Finch (*Taeniopygia castanotis*).

In addition a Kingfisher, Babbler, another Blue-Wren, Crow, and Magpie are present, but the species are not definitely identified.

*Bird Notes.*

Crested Pigeon (*Ocyphaps lophotes*); Southern Stone Curlew (*Burhinus magnirostris*); Bustard (*Eupodotis australis*); Grey Teal (*Querquedula gibberifrons*); Wedge-tailed Eagle (*Uroaetus audax*); Black-shouldered Kite (*Elanus axillaris*); Brown Hawk (*Falco berigora*); Nankeen Kestrel (*Falco cenchroides*); White Cockatoo (*Kakatoe galerita*) (?); Galah (*K. roseicapilla*); Cockatoo Parrot (*Leptolophus hollandicus*); Port Lincoln Parrot (*Barnardius zonarius*); Mulga Parrot (*Psephotus varius*); Budgerygah (*Melopsittacus undulatus*); Owlet Nightjar (*Aegotheles cristata*); Bee-eater (*Merops ornatus*); Welcome Swallow (*Hirundo neoxena*); Tree-Martin (*Hylochelidon nigricans*); Fairy Martin (*H. ariel*); Magpie-Lark (*Grallina cyanoleuca*); Crested Bellbird (*Oreoica gutturalis*); Black-faced Cuckoo-Shrike (*Coracina novae-hollandiae*); Brown Songlark (*Cinclorhamphus cruralis*); Blue and White Wren (*Malurus cyanotus*); White-plumed Honeyeater (*Meliphaga penicillata*); Yellow-throated Miner (*Myzantha flavigula*); Spiny-cheeked Honeyeater (*Acanthagenys rufogularis*); Chestnut-eared Finch (*Taeniopygia castanotis*).

In addition a Kingfisher, Babbler, another Blue-Wren, Crow, and Magpie are present, but the species are not definitely identified.